



Incorporating Short Data into Large Mixed-Frequency VARs for Regional Nowcasting

Gary Koop, Stuart McIntyre, James Mitchell, Aubrey
Poon and Ping Wu

ESCoE Discussion Paper 2023-07

May 2023

ISSN 2515-4664

DISCUSSION PAPER

Incorporating short data into large mixed-frequency VARs for regional nowcasting
Gary Koop, Stuart McIntyre, James Mitchell, Aubrey Poon and Ping Wu
ESCoE Discussion Paper No. 2023-07
May 2023

Abstract

Interest in regional economic issues coupled with advances in administrative data is driving the creation of new regional economic data. Many of these data series could be useful for nowcasting regional economic activity, but they suffer from a short (albeit constantly expanding) time series which makes incorporating them into nowcasting models problematic. Regional nowcasting is already challenging because the release delay on regional data tends to be greater than that at the national level, and “short” data imply a “ragged edge” at both the beginning and the end of regional data sets, which adds a further complication. In this paper, via an application to the UK, we develop methods to include a wide range of short data into a regional mixed-frequency VAR model. These short data include hitherto unexploited regional VAT turnover data. We address the problem of the ragged edge at both the beginning and end of our sample by estimating regional factors using different missing data algorithms that we then incorporate into our mixed-frequency VAR model. We find that nowcasts of regional output growth are generally improved when we condition them on the factors, but only when the regional nowcasts are produced before the national (UK-wide) output growth data are published.

Keywords: Regional data, Mixed-frequency data, Missing data, Nowcasting, Factors, Bayesian methods, Real-time data, Vector autoregressions

JEL classification: C32, C53, E37

Stuart McIntyre, University of Strathclyde
s.mcintyre@strath.ac.uk

Published by:
Economic Statistics Centre of Excellence
King's College London
Strand
London
WC2R 2LS
United Kingdom
www.escoe.ac.uk

ESCoE Discussion Papers describe research in progress by the author(s) and are published to elicit comments and to further debate. Any views expressed are solely those of the author(s) and so cannot be taken to represent those of the Economic Statistics Centre of Excellence (ESCoE), its partner institutions or the Office for National Statistics (ONS).

© Gary Koop, Stuart McIntyre, James Mitchell, Aubrey Poon and Ping Wu

Incorporating Short Data into Large Mixed-Frequency VARs for Regional Nowcasting*

Gary Koop,[†] Stuart McIntyre,[‡] James Mitchell,[§] Aubrey Poon,[¶] and Ping Wu^{||}

May 2023

Abstract

Interest in regional economic issues coupled with advances in administrative data is driving the creation of new regional economic data. Many of these data series could be useful for nowcasting regional economic activity, but they suffer from a short (albeit constantly expanding) time series which makes incorporating them into nowcasting models problematic. Regional nowcasting is already challenging because the release delay on regional data tends to be greater than that at the national level, and “short” data imply a “ragged edge” at both the beginning and the end of regional data sets, which adds a further complication. In this paper, via an application to the UK, we develop methods to include a wide range of short data into a regional mixed-frequency VAR model. These short data include hitherto unexploited regional VAT turnover data. We address the problem of the ragged edge at both the beginning and end of our sample by estimating regional factors using different missing data algorithms that we then incorporate into our mixed-frequency VAR model. We find that nowcasts of regional output growth are generally improved when we condition them on the factors, but only when the regional nowcasts are produced before the national (UK-wide) output growth data are published.

Keywords: Regional data, Mixed-frequency data, Missing data, Nowcasting, Factors, Bayesian methods, Real-time data, Vector autoregressions

JEL Codes: C32, C53, E37

*This research has been funded by the ONS as part of the research program of the Economic Statistics Centre of Excellence (ESCoE). The views expressed herein are those of the authors and not necessarily those of the Federal Reserve Bank of Cleveland or the Federal Reserve System.

[†]University of Strathclyde; Economic Statistics Centre of Excellence (gary.koop@strath.ac.uk)

[‡]University of Strathclyde; Economic Statistics Centre of Excellence (s.mcintyre@strath.ac.uk)

[§]Federal Reserve Bank of Cleveland; Economic Statistics Centre of Excellence (james.mitchell@clev.frb.org)

[¶]Orebro University; Economic Statistics Centre of Excellence (aubrey.poon@oru.se)

^{||}University of Strathclyde; Economic Statistics Centre of Excellence (ping.wu@strath.ac.uk)

1 Introduction

“Official” sub-national data, as published by national statistical institutes, tend to be available at a lower frequency and published more slowly than data for the nation as a whole. A case in point, and the motivating empirical example in this paper, is regional data for gross value added (GVA) for the UK regions.¹ While the Office for National Statistics (ONS) has long produced GVA data for the UK regions, until 2012 these data were produced at an annual frequency only and with a release delay of approximately one year.² In 2012, with the development of the quarterly regional GDP data set, ONS production was sped up. Regional GVA then became available at a quarterly frequency and with a shorter, but still substantial, release delay of approximately six months. This situation is not unusual at a regional level.³ There therefore remains a need, especially to support policy and business decisions made in real time, to produce more timely regional estimates – so-called “nowcasts.”

The motivation for this paper is the observation that recent years have seen increasing availability, from both private and public sources, of higher-frequency and more timely data, capturing various aspects of economic activity, at the regional and national levels. Ideally, when nowcasting, one would condition on all of these indicators - and let the data (the fit of the model) determine which indicators are most useful. But these new indicators are often available with only a limited historical time series. This impedes their inclusion in traditional nowcasting models. Table 1 illustrates the issue in the context of our UK application (but we emphasize that this issue arises for other variables and in other countries). From Table 1, we see that many regional indicators are available only from the 1990s onward; e.g., Labour Force Survey data produced by the ONS, house price indices, and the Purchasing Managers’ Index (PMI). But perhaps of most interest are some of the recently released data series based on administrative data that have not been previously used for regional nowcasting in the UK.

These administrative data include payroll employment information derived from the “pay as you earn” (PAYE) tax system, which are available back to July 2014 only, and value added tax (VAT) turnover data, which are available back to 2012. There are good reasons to think that these new “short” indicators should be useful in regional nowcasting. Especially so when, like the PAYE and VAT data, they are derived from administrative data and, therefore, reflect the universe of individuals/firms covered by particular taxes. This provides a significant improvement upon survey-based measures of activity. The payroll employment data also benefit from being high frequency (monthly) and very timely (released within two weeks of the end of the month). The VAT data, while quarterly and released with a delay of around 5 months (this reflects the time taken to produce consistent sub-national aggregates from the underlying firm-level VAT returns), are a key input into the ONS’s

¹Real GVA and real GDP are closely related concepts. GVA is in basic prices, while GDP is in market prices (i.e., GVA plus taxes (less subsidies) on products equals GDP).

²In this paper, we consider the 12 ITL1 regions, excluding the UK continental shelf. These 12 regions/devolved nations comprise North East England, North West England, Yorkshire and the Humber, East Midlands, West Midlands, East of England, London, South East England, South West England, Wales, Scotland, and Northern Ireland.

³In the European Union, for example, regional output data are available only annually (at the ITL2 level) and with a release delay of more than a year, while in the US state-level GDP data are available quarterly, but with a release delay of around 3 months - in contrast to data for the US as a whole, which are released with a delay of 4 weeks.

production of regional GDP data themselves. We should therefore expect VAT data to provide useful information when nowcasting regional GVA growth. But the short and differing lengths of these new data mean that questions arise over how to include them as indicators in a nowcasting model.

Researchers wishing to update their nowcasts using the latest available information are used to addressing what is commonly called, since Wallis (1986), the “ragged edge” problem. Since different variables are released with different delays, variables with longer delays will have missing values at the end of the sample, whereas variables with shorter release delays will not. This leads to a ragged edge at the bottom of the spreadsheet. Here we have this issue, but we also have a similar problem occurring at the beginning of the sample. The question of how to address the ragged edge at *both* the beginning and the end of the sample in the context of a regional nowcasting model is thus the focus of this paper. Given that the regional GVA data themselves are available in some form back to 1966, there is potentially a large loss in estimation accuracy and efficiency if the model is simply estimated only over the common sample; in the context of Table 1, this would be from the 1990s onward. It is preferable to consider all available data across both the time-series and the cross-sectional dimensions. However, data with a short time span are difficult to incorporate into conventional time-series econometric models, which often work best with long samples.

We address this challenge by developing methods to incorporate these “short” data, and potentially a large number of “short” indicators, into mixed-frequency regional nowcasting models. The specific class of model we consider is a mixed-frequency vector autoregression (MF-VAR). MF-VARs are a popular nowcasting tool; for example, see Brave et al. (2019); Eraker et al. (2014); Schorfheide and Song (2015); McCracken et al. (2021). Koop et al. (2020b,c) extended MF-VAR models to the regional case by incorporating an additional measurement equation reflecting the fact that national GDP is the aggregation of regional GDP: they call this the “cross-sectional constraint.” This paper sets out how to use this class of MF-VAR models when nowcasting using short data of the type seen in Table 1. The proposed model involves a large number of regional variables, some with much shorter time spans than others, and using these to construct factors for each region using factor extraction methods. These factors, summarizing the information in the regional predictors, are then incorporated into the MF-VAR. This leads to a mixed-frequency factor-augmented VAR (MF-FAVAR). At the beginning of the sample, the regional factors reflect information in only a few regional indicators. But as time passes and more variables become available, these data are also included in the factors.

Our point of departure is to consider how best to construct these regional factors from the mixed-frequency and ragged edge data shown in Table 1. Traditionally, missing data problems in factor-based macroeconomic analysis have been addressed using expectation maximization principal components analysis (EMPCA); see Stock and Watson (2002). More recently, new approaches have been suggested, including the tall wide (TW) algorithm of Bai and Ng (2021) and the tall project (TP) algorithm of Cahan et al. (2023). These are all generic factor estimation algorithms to handle missing data. We consider the relative performance of these different algorithms when dealing with our specific missing data configuration - the ragged edge at the beginning and end of the sample - and how to incorporate these factors into the MF-FAVAR.

Importantly, the model developed has wide applicability for regional nowcasting applications. It can be used to accommodate other evolving and ragged edge data features that might be evident for other variables and in other countries. This includes more (or less) timely and higher (lower) frequency regional output (GVA) data. This model overcomes a major barrier to the more widespread use of new innovative data sources for regional nowcasting. Given the trend of increasing data availability at a regional level, and interest from policymakers in improving the spatial granularity of economic statistics, the development of a regional nowcasting model that is robust to different release delays, time-series lengths, and data frequencies, represents a valuable addition to the toolkit available to researchers. Via a UK application, we explore how our model performs in a pseudo-real-time nowcasting exercise and we test whether including these additional predictors improves the accuracy of our nowcasts of UK regional GVA.

The remainder of this paper is structured as follows. In Section 2 we describe in more detail the data we use in our empirical work. In Section 3 we set out the econometric methods used for nowcasting, including factor analysis of missing data. We explore the properties of three algorithms for the estimation of factors with ragged edge samples via a set of Monte Carlo exercises. Section 4 reports our empirical results, including a comparison of the factors produced across different methods and their performance in a pseudo-real-time nowcasting exercise. Section 5 concludes.

2 The Data

In this section, we set out the data used in our model. We do this in two parts. First, we summarize the regional output growth data. Second, we describe the predictors that we incorporate into our model, via the calculation of regional factors, to nowcast quarterly regional output growth.

2.1 Regional Output Growth Data

We combine “official” annual regional real GVA growth data produced by the ONS since 1966 with quarterly regional GVA growth data once they become available. For the English regions and Wales, the ONS quarterly data date back to 2012Q1; Scottish government data date back to 1998Q1; and Northern Ireland Statistics and Research Agency data go back to 2006Q1. Pre-1998, the annual GVA data are available only in nominal terms, and we follow previous research and construct a historical regional database for the UK by using a UK GDP deflator. For more details on the construction of this regional database, see Koop et al. (2020a,b,c, Forthcoming). In the absence of real-time data vintages for the regional GVA data, we consider only the latest-vintage GVA data.⁴

2.2 Regional Indicators: Short and Long Data

An increasing volume of regional economic data, from both official and unofficial sources, is becoming available. While we cannot claim that the data set we use in this paper represents complete coverage

⁴In general, the release delay for the regional GVA data is 6 months, but the delay for the most recent data has been longer, meaning that at the time of writing the latest vintage is 2021Q4.

of the regional data available, it does represent a data set with features typical of regional economic data and it has coverage across many different types of economic data. For use in nowcasting, any data have to be released on a more timely basis than the target variable (in our case, regional GVA). This rules out annual indicators and any quarterly indicators that are released with too long a delay after the quarter to which they relate.⁵ A number of other indicators exist, but we do not have access to them for a variety of reasons.⁶ However, we should stress that the model we develop in this paper is capable of incorporating such data should they become available.

Having set out the criteria for data inclusion, Table 1 summarizes the indicators that we ultimately include in the model. These indicators cover a range of time periods, with some measures having relatively short time spans (for example, payroll employment data that only go back to 2014), while others (for example, the house price information) cover the entire sample period. We also observe a range of release delays, with some indicators being released very quickly after the end of the reference period, and others being only slightly more timely than regional GDP itself. The indicators included in this model can be grouped into three main categories: measures of output, labor market data, and housing market indicators. Measures of output include data covering: construction sector output, retail sales, trade data, port traffic, tourism stays and spending, and business demographic information. As discussed, we also included output indices only available for Scotland and Northern Ireland (for example, the Northern Ireland Index of Services and the Northern Ireland Index of Production). At a quarterly frequency, there is also one survey of the business outlook, the CBI business optimism index available for all ITL1 regions, and a Scottish consumer sentiment indicator. A key monthly indicator of business activity is the PMI survey measure. This comprises separate indicators of activity that were each included separately: new business, outstanding business, charges, prices, employment, and future orders. For Scotland, there is also a monthly GDP measure.

This paper is the first to utilize VAT turnover aggregates at the regional level, provided to us by the ONS, in a pseudo-real-time nowcasting exercise. These data are disaggregated to the ITL3 (formerly NUTS3) regions of the UK, and are available on a quarterly basis from 2012Q1. For each ITL1 region, we use VAT data for that ITL1 region, as well as each of the ITL2 and ITL3 subregions within it, as separate variables used in the construction of the regional factors. Vintage data are available only from 2019Q4 (which, combined with the previously mentioned lack of real-time data for regional output growth, also explains our need to undertake a pseudo-real-time exercise). The typical publication lag of these data is 5 months after the end of the reference quarter. These data are produced by the ONS (although they are not typically available to researchers outside of the ONS). The data are aggregations of individual VAT returns from firms, which are cleaned by ONS to address any anomalies in the completion of these forms by businesses and the declared turnover data assigned to a quarter and a geographic location (reflecting an allocation to each ITL1/2/3 level). The

⁵For example, in the UK there are local-level data on lending to small and medium-size enterprises and also mortgage lending, but neither is more timely than our target measure of regional output.

⁶For example, the need for a commercial subscription (e.g., GfK consumer confidence data), the lack of data in aggregate form (e.g., ONS's Monthly Business Survey microdata and credit/debit card transactions data), or the data being privately held.

Table 1: Regional Indicators: Short and Long Data

Variable	Tab	Description	Frequency	Geographic Coverage (National - N; Regional-R)	Time period	Typical release delay
1	UK Quarterly HPI	Nationwide Building Society House Price Index	Q	N	Q1 1967 -	1 week
2	UKEMP	16 - 64 Employment Rate Labour Force Survey	Q	N	Q2 1992 - Q4 2021	6 weeks
3	UKUnEMP	16+ Unemployment Rate Labour Force Survey	Q	N	Q2 1992 - Q4 2021	6 weeks
4	UKMonthlyHousePrice	UK Government UK House Price Index	M	N	April 1968 -	6 weeks
5	RegCBI	CBI Business Optimism index	Q	R	Q2 1958 - Q1 2021	4 weeks
6	BusinessBirths	Business Births Geography Counts	Q	R	Q1 2017 - Q1 2022	1 month
7	ConstructionOutput_TNH	Construction Output - Total new housing	Q	R (ex. NI)	Q1 1980 - Q4 2021	6 weeks
8	ConstructionOutput_ANW	Construction Output - All New Work	Q	R (ex. NI)	Q1 1980 - Q4 2021	6 weeks
9	ConstructionOutput_AW	Construction Output - All Work	Q	R (ex. NI)	Q1 1980 - Q4 2021	6 weeks
10	Employment	16 - 64 Employment Rate Labour Force Survey	Q	R	Q2 1992 - Q4 2021	6 weeks
11	Unemployment	16+ Unemployment Rate Labour Force Survey	Q	R	Q2 1992 - Q4 2021	6 weeks
12	PublicEMP	Public sector employment	Q	R	Q1 2008 - Q4 2021	3 months
13	WorkforceJobs	Workforce jobs by region and industry	Q	R	Q1 1996 - Q1 2022	3 months
14	Exports	Value of Exports	Q	R	Q1 2018 - Q3 2021	4 months
15	Imports	Value of Imports	Q	R	Q1 2018 - Q3 2021	4 months
16	RegCC	Claimant count rate	M	R	Apr 1974 - Jan 2022	2 weeks
17	PayrollEMP	Payroll employment	M	R	July 2014 - Jan 2022	2 weeks
18	PayrollPayMedian	Median payroll pay	M	R	July 2014 - Jan 2022	2 weeks
19	HousePrice	House Price Index	M	R	Jan 1995 - Mar 2022 (most regions, some start later)	6 weeks
20	PMIactivity	PMI activity measure (headline)	M	R	Jan 1997 - Jan 2022 (most regions, some start later)	2 weeks
21	PMInewbus	New business measure	M	R	Jan 1997 - Jan 2022 (most regions, some start later)	2 weeks
22	PMIoutbus	Outstanding business	M	R	Nov 1999 - Jan 2022 (most regions, some start later)	2 weeks
23	PMIcharges	Charges	M	R	Nov 1999 - Jan 2022 (most regions, some start later)	2 weeks
24	PMIprices	Prices	M	R	Jan 1997 - Jan 2022 (most regions, some start later)	2 weeks
25	PMIemploy	Employment	M	R	Jan 1997 - Jan 2022 (most regions, some start later)	2 weeks
26	PMIfuture	Future orders	M	R	Jan 1997 - Jan 2022 (most regions, some start later)	2 weeks
27	HousingRental	Private Housing Rental Prices	M	R	Jan 2012 - Jan 2022 (most regions, some start later)	2 weeks
28	OvernightVisits	Number of overnight visits to the regions of the UK by area of residence	Q	R	Jan 2005 - Apr 2022 (most regions, some start later)	3 weeks
29	Spending	Spending by overseas residents in regions of the UK by area of residence	Q	R (selected regions)	Q1 2017 - Q3 2021	5 months
30	IncorporatedCompanies	Number of companies on the register, newly incorporated companies, and removals from the register.	Q	R (selected regions)	Q1 2009 - Q3 2021	5 months
31	Scot_GDP	Scottish monthly GDP	M	R (Scot only)	Jan 2011 - Q1 2022	1 month
32	Scot_LabourProductivity	Scottish Labour productivity	Q	R (Scot only)	Jan 2010 - Mar 2022	2 months
33	Scot_RetailSales	Retail sales index for Scotland	Q	R (Scot only)	Q1 1998 - Q4 2019	5 months
34	Scot_CSI	Scottish Consumer Sentiment Indicator	Q	R (Scot only)	Q1 2008 - Q1 2020	1 month
35	NI_IOS	Northern Ireland Index of Services	Q	R (NI only)	Q2 2013 - Q1 2022	1 month
36	NI_IOP	Northern Ireland Index of Production	Q	R (NI only)	Q1 2005 - Q4 2021	3 months
37	NI_RSI	Northern Ireland Retail Sales Index	Q	R (NI only)	Q1 2005 - Q4 2021	3 months
38	NI_Ports Traffic	Northern Ireland Ports Traffic	Q	R (NI only)	Q1 2014 - Q4 2021	3 months
39	NI_ConstructionOutput	Construction output in Northern Ireland	Q	R (NI only)	Q1 2009 - Q4 2021	4 months
40	VAT	VAT Turnover by ITL1/2/3 Regions	Q	R	Q1 2013 - Q4 2021	3 months
					Q1 2012 - Q4 2021	5 months

issues involved in this data work by ONS are significant and challenging; see, for example, Labonne and Weale (2020).

Labour Force Survey (LFS) data for the ITL1 regions of the UK are also incorporated. They are available with a release delay of around 6 weeks after the end of the quarter to which they relate. These data are released on a rolling 3-month basis and are updated each month. The LFS provides a range of measures of labor market activity, from which we select headline employment and unemployment, and public-sector employment. There are a small number of monthly labor market measures. These include data from the social security system (for example, the claimant count rate), as well as the HMRC real-time information data from the PAYE system on payroll employment and pay. Given the important role of the housing market in the economy (see Leamer (2007)), we also include two measures of changes in house prices. The first of these is monthly house price index data for each ITL1 region published by the UK government, as well as the quarterly UK house price index published by the Nationwide Building Society (which has the advantage of being more timely than the official data series). We also include information on rental prices for the private rental market.

3 Econometric Methods

3.1 Notation and Data Availability

We begin by describing some variable definitions, relationships, and notational conventions used in this paper.

- $t = 1, \dots, T$ runs at a *quarterly* frequency.
- $r = 1, \dots, R$ denotes the R regions in the UK.
- Y_t^{UK} is GVA for the UK in quarter t .
- $y_t^{UK} = \log(Y_t^{UK}) - \log(Y_{t-1}^{UK})$ is the quarterly change (log difference) in GVA in the UK.
- Y_t^r is GVA for region r in quarter t . It is not observed before 2012 except for Scotland and Northern Ireland, where it is not observed before 1998 and 2006, respectively.
- $Y_t^{r,A} = Y_t^r + Y_{t-1}^r + Y_{t-2}^r + Y_{t-3}^r$ is annual GVA for region r . It is observed in quarter 4 of each year, but not in other quarters.
- $y_t^{r,A} = \log(Y_t^{r,A}) - \log(Y_{t-4}^{r,A})$ is annual GVA growth in region r . It is observed, but only in quarter 4 of each year. $\mathbf{y}_t^A = (y_t^{1,A}, \dots, y_t^{R,A})'$ is the vector of annual GVA growth rates for the R regions.
- $y_t^r = \log(Y_t^r) - \log(Y_{t-1}^r)$ is the quarterly change in GVA in region r . It is not observed before 2012 except for Scotland and Northern Ireland, where it is not observed before 1998 in Scotland and 2006 in Northern Ireland. $\mathbf{y}_t^Q = (y_t^1, \dots, y_t^R)'$ is the vector of quarterly GVA growth rates for the R regions.

- The link between the quarterly regional growth rates and their annual counterpart is referred to as the inter-temporal restriction and takes the form:

$$y_t^{r,A} = \frac{1}{4}y_t^r + \frac{1}{2}y_{t-1}^r + \frac{3}{4}y_{t-2}^r + y_{t-3}^r + \frac{3}{4}y_{t-4}^r + \frac{1}{2}y_{t-5}^r + \frac{1}{4}y_{t-6}^r. \quad (1)$$

This restriction is not imposed in periods where quarterly regional GVA growth data are available.

- The link between the regional growth rates and the UK counterpart is referred to as the cross-sectional restriction and takes the form:⁷

$$y_t^{UK} \approx \frac{1}{R} \sum_{r=1}^R y_t^r. \quad (2)$$

- \mathbf{Z}_t^r is a vector containing k_r quarterly variables for region r . These are the “short” data that start at differing times as described in Section 2.

3.2 The MF-FAVAR

To explain the structure of our MF-FAVAR, we begin with a structural VAR that relates a vector of N dependent variables, \mathbf{y}_t to lags of the dependent variables:

$$\mathbf{B}\mathbf{y}_t = \mathbf{A}\mathbf{x}_t + \boldsymbol{\varepsilon}_t \quad (3)$$

where \mathbf{x}_t is a vector containing p lags of \mathbf{y}_t . The errors, $\boldsymbol{\varepsilon}_t$, are assumed to be $\mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma})$, where $\boldsymbol{\Sigma}$ is a diagonal matrix and \mathbf{B} is lower triangular with ones on the diagonal.⁸

In a conventional VAR, all of the dependent variables are simply observed variables. Note that \mathbf{A} is an $N \times Np$ matrix and, thus, there are pN^2 VAR coefficients to be estimated. If N and/or p is large, VARs can be seriously over-parameterized. We investigated lag lengths of up to $p = 7$ (and our results are robust, so our main results set $p = 1$). Thus, our \mathbf{A} matrix is big. Accordingly, we use Bayesian estimation methods that allow for prior shrinkage.

The MF-VAR is a VAR where \mathbf{y}_t is no longer simply a set of *observed* variables. Instead, some of the elements in \mathbf{y}_t are unobserved or latent variables. In particular, there are the unobserved high-frequency values of the low-frequency variables: the objects we wish to estimate. The latter are linked to the former via the inter-temporal restriction. We set $\mathbf{y}_t = \left(y_t^{UK}, \mathbf{y}_t^{Q'} \right)$, where \mathbf{y}_t^Q contains the quarterly regional growth rates that we were seeking to estimate.

⁷The cross-sectional restriction given here assumes growth rates are modeled as log differences; see Koop et al. (2020c) for the derivation of this restriction. As discussed in this paper, the constraint is approximate to reflect both the logarithmic approximation and the fact that regional output does not exactly sum to UK output because of the UK continental shelf.

⁸Working in this structural VAR form does not restrict the reduced-form error covariance matrix and greatly simplifies computation, since estimation can proceed one equation at a time.

The MF-VAR can be set up as a state space model, where the state equations are given by the VAR and the measurement equations are the inter-temporal and cross-sectional restrictions. Bayesian methods exist for posterior and predictive inference in such state space models. For details, see Koop et al. (2020c,b). We use variational Bayes (VB) methods instead of Markov chain Monte Carlo (MCMC) estimation, because of their computational advantages. VB methods for the MF-VAR are developed in Gefang et al. (2020) and used by Koop et al. (Forthcoming). In this paper, we also use a computationally more efficient precision-based approach to estimate the states, instead of the Kalman filter; see Chan et al. (2021).

The MF-FAVAR we use is an MF-VAR, but with one final re-definition of \mathbf{y}_t . In particular, it sets $\mathbf{y}_t = (y_t^{UK}, y_t^{Q'}, \mathbf{f}_t')$, where $\mathbf{f}_t = (\mathbf{f}_t^1, \dots, \mathbf{f}_t^R)'$ is a vector of regional factors. It is possible to have more than one factor for each region and, accordingly, each of the \mathbf{f}_t^r is a vector of n_f factors constructed using \mathbf{Z}_t^r for $r = 1, \dots, R$. These factors are observed at a quarterly frequency and, thus, can be treated as additional high-frequency observed variables in the MF-VAR. Bai and Ng (2006) establish the conditions under which the estimated factors can be treated as known in estimation when doing Principal components analysis (PCA). Since all of our methods involve PCA (but differ in their treatment of missing values) the results of Bai and Ng (2006) suggest that little is lost by treating the estimated factors as known.

It is worth noting that our MF-FAVAR is of very high dimension, including at a minimum $N = R \times n_f + 1$ dependent variables. This high dimension arises since, for each region, we are including regional output growth plus n_f factors and for the UK we include at least GVA growth. Even if we only include one factor for each of the 12 regions and no additional UK-wide variables in the model, we end up with a VAR of dimension $N = 25$, which is already quite large. In practice, the need to include more than one factor and/or additional UK variables means that most of our models are of a much higher dimension.

We partially surmount the over-parameterization problem by working with a restricted version of the MF-FAVAR. This imposes the restriction that the equation for GVA growth for a particular region depends only on the regional factor for that region (as well as UK variables and lags of GVA growth for that region). In other words, each regional factor is specific to a region and does not appear in the equation for other regions. As a robustness check, we also estimated the unrestricted version of the model; see the online appendix.

We also use Bayesian prior shrinkage as a way of avoiding over-parameterization concerns. There exist a range of VAR priors and any of them could be used. In this paper, we use the popular adaptive Lasso (AL). This is a global-shrinkage prior that automatically chooses which coefficients should be shrunk to zero; see Zou (2006) for details. We have experimented with two versions of the AL prior. One of these uses the AL on all the coefficients in the model (except for error variances, which are the diagonal elements of Σ for which we use relatively non-informative inverse-Gamma priors). In the other version we use the AL for all coefficients except for the error covariances (i.e., the parameters controlling the contemporaneous relationships between the variables in the model). For these, we adopt the asymmetric conjugate prior (ACP) of Chan (2022), which might be expected to have good

properties for these parameters. We name our two priors AL and AL-ACP, respectively. In practice we find that AL-ACP leads to slightly better forecasts and hence the results presented in the main body of the paper are for AL-ACP. In the online appendix, we present the AL results.

In summary, our algorithm begins by constructing the factors, using methods described in the next sub-section, and then includes them in the MF-FAVAR. We then use Bayesian methods to construct nowcasts of quarterly GVA growth for the UK regions.

3.3 Constructing Regional Factors with Short Data

In this section, we describe three methods for constructing factors from \mathbf{Z}_t^r , which, as seen in Table 1, is a matrix characterized by missing data at the beginning and end of the sample. These algorithms are denoted by the abbreviations EMPCA, TW, and TP. We now provide a brief description of each algorithm in turn, along with a summary of its properties. Full details are provided in Stock and Watson (2002) for EMPCA, in Bai and Ng (2021) for TW, and in Cahan et al. (2023) for TP.

All of these approaches are algorithms for filling in missing data where various patterns of missingness are possible. In the present paper, as discussed, we are especially interested in one particular pattern of missingness: the ragged edge at the beginning of the sample due to the short nature of many of the \mathbf{Z}_t^r indicators. In order to explore the relative performance of the three algorithms across data sets with “short” data, we simulate data sets with different properties (including sample size, number of variables, and degree of missingness) in a set of Monte Carlo experiments summarized in Section 3.3.4 below. We first summarize the three factor estimation methods.

3.3.1 EMPCA

PCA is a popular method for estimating factors. It is a non-parametric method. The advantage of this is that it is less liable to specification error than parametric approaches. But this can also be a disadvantage, as sensible parametric assumptions can improve estimation accuracy (for example, in a macroeconomic context, factors might be expected to exhibit autoregressive behavior and a parametric model that allows for such behavior could improve estimation accuracy). As discussed, for example in Section 2.3 of Stock and Watson (2016), PCA exploits correlations across variables to produce estimates of the factors.

When data are missing, Stock and Watson (2002) combine PCA with methods to fill in missing data using an expectations-maximization (EM) algorithm, leading to EMPCA. In our context, each regional factor, \mathbf{f}_t^r , is calculated using \mathbf{Z}_t^r which contains missing values. Let $\hat{\mathbf{Z}}_t^r$ be a version of \mathbf{Z}_t^r with these missing values replaced by estimates and $\hat{\mathbf{f}}_t^r$ denote estimates of the factors. EMPCA is an iterative algorithm that uses PCA on $\hat{\mathbf{Z}}_t^r$ to produce $\hat{\mathbf{f}}_t^r$. Estimates of the missing values are produced as fitted values from a regression of $\hat{\mathbf{Z}}_t^r$ on $\hat{\mathbf{f}}_t^r$.

3.3.2 TW

The tall wide (TW) algorithm of Bai and Ng (2021) is not an iterative algorithm and, thus, does not suffer from the computational issues that can afflict EMPCA. To understand TW, consider the $T \times k_r$ matrix of data for a specific region, \mathbf{Z}_t^r . Each column contains all the data for a variable. Some of the columns will not have any missing values for any time periods. These columns form what is called the “tall” block; first undertake PCA using this block, produce factors $\hat{\mathbf{f}}_t^{r,Tall}$ and factor loadings $\hat{\mathbf{A}}^{r,Tall}$. Next consider the rows of \mathbf{Z}_t^r . Some of these rows will not have missing values and these rows will form the “wide” block. PCA can be applied to this wide block to produce factors $\hat{\mathbf{f}}_t^{r,Wide}$ and factor loadings $\hat{\mathbf{A}}^{r,Wide}$. The TW algorithm is a simple method for combining $\hat{\mathbf{f}}_t^{r,Tall}$, $\hat{\mathbf{A}}^{r,Tall}$, $\hat{\mathbf{f}}_t^{r,Wide}$, and $\hat{\mathbf{A}}^{r,Wide}$ in an optimal way (using least squares regression methods) to produce a single regional factor.

Bai and Ng (2021) show the TW algorithm to have desirable asymptotic properties. These properties depend on the number of columns/rows in the tall/wide blocks. Note that if the tall block is very narrow (that is, few variables have data available for the full sample), then $\hat{\mathbf{f}}_t^{r,Tall}$ will be based on few variables and may be a poor estimate of the regional factor(s). Similarly, if the wide block is thin, then $\hat{\mathbf{f}}_t^{r,Wide}$ will only be available for a few observations and may produce poor estimates. In our regional nowcasting context, suppose, for instance, that the variable with the latest start date begins in 2010. In this case, $\hat{\mathbf{f}}_t^{r,Wide}$ would only be available for 2010 onward. Furthermore, all the observations pre-2010 will be discarded other than observations for variables in the tall block. We might therefore expect the TW algorithm to be sensitive to the choice of variables and that it would not work well if one (or a few) of the variables is available for a short period of time.

3.3.3 TP

The tall project (TP) algorithm of Cahan et al. (2023) is similar to TW in that the tall block plays a key role and $\hat{\mathbf{f}}_t^{r,Tall}$ and $\hat{\mathbf{A}}^{r,Tall}$ are key ingredients in the estimated regional factor(s). However, it surmounts the problem noted previously that occurs with the TW algorithm when one of the variables has a very short time and, thus, the wide block is thin. It does so by using auxiliary regressions for the observed values of each individual variable (other than those in the tall block) on the tall block factors. The auxiliary regression for variable i can be used to fill in the missing values for variable i , thus leading to $\hat{\mathbf{Z}}_t^r$ that does not have missing values. The regional factor is estimated using PCA on $\hat{\mathbf{Z}}_t^r$. With this algorithm, it is possible to iterate, but Cahan et al. (2023) show that asymptotically this is not necessary. In this paper, we do not iterate.

3.3.4 Monte Carlo Evaluation of the Three Factor Algorithms

This section summarizes the results from a set of Monte Carlo experiments designed to evaluate the EMPCA, TW, and TP algorithms when applied to data sets with ragged edges, to varying degrees, at the beginning of the sample. Full details of the data-generating process and the simulation exercise are in online Appendix A.

EMPCA is an iterative algorithm that raises two computational issues: it is fundamentally slower than non-iterative methods and can fail to converge. TW and TP are simpler algorithms, fast, and not subject to concerns about convergence. However, they are likely to be more sensitive to the number of variables without missing observations (that is, the size of the tall block). In this paper, where our application involves a substantial ragged edge at the beginning of the sample and our tall block potentially contains only a small number of variables, it is possible that these properties of the TW and TP algorithms will make it worthwhile to take on the larger computational burden of EMPCA. But the choice between the three algorithms is fundamentally an empirical issue.

To assess the precision of the estimates of the factors from the three algorithms, we conduct a set of Monte Carlo experiments. Data are generated from the factor model used in Bańbura and Modugno (2014). We generate data for different sample lengths T , different sizes of the cross-sectional panel n , and different values for τ (which governs the degree of cross-correlation of the idiosyncratic component). Then we leave the first two simulated variables as complete: this is our tall block. We let two variables remain complete because in our regional nowcasting application we have two indicators with no missing data. For the remaining $(n - 2)$ variables, we set a certain fraction of the data as missing. Given our interest in the ragged edge at the beginning of the sample, we place these missing data points at the beginning of the sample. We consider cases of 0, 20, 40, 60, and 80 percent of missing data. Then, using the simulated data, we estimate factors using the three algorithms.

To evaluate the precision of the factor estimates, we follow Bańbura and Modugno (2014) and compute the trace R^2 of the regression of the estimated factors on the true ones. Table A1 in Appendix A reports average trace statistics over 500 Monte Carlo replications for EMPCA, and Table A2 reports the statistics for TW and TP. From these tables we see that the three algorithms have similar estimation accuracy. The estimates are less precise for small sample lengths ($T = 50$ versus $T = 200$), small cross-sections ($n = 12$ versus $n = 102$), a mis-specified model ($\tau > 0$ versus $\tau = 0$) in small samples, and a large fraction of missing data. Table A1 also shows that the EMPCA algorithm is slower than the other two approaches. But the additional computational burden is small. More significantly, however, there were some instances of convergence failures. This included samples where $T = 200$, $n = 50$, and 60 percent of the sample is missing: this matches the features of our regional nowcasting data set (seen in Table 1). Thus we suspect that there might be convergence problems when using EMPCA in our application.

4 Regional Nowcasting with the MF-FAVAR

4.1 Design of the Nowcasting Exercise and Specification Choices

In our regional nowcasting exercise, we will compare different versions of our MF-FAVAR to an MF-VAR that is identical (that is, same prior, same lag length choice, etc.), except that it does not include the regional factors extracted from the short (and longer) data summarized in Table 1. The MF-VAR is our “benchmark” model. In terms of the indicators in the models, all include GVA growth for the 12 regions plus the UK as a whole, along with four quarterly UK macroeconomic predictors

(inflation, interest rates, the exchange rate, and oil price inflation). Thus, our smallest model, the MF-VAR, includes 17 variables. Then in the MF-FAVARs we add a set of regional factors. Note that the factor for region r is based only on data from that region (i.e. \mathbf{Z}_t^r contains data specific to region r , including sub-regional VAT data for region r). The variables are given in Table 1. The number of variables used to construct the factors will differ across regions due to data availability and the fact that the number of sub-regions varies across regions.

The MF-VARs differ in four ways:

1. The way the factors are calculated (EMPCA, TW, or TP);
2. The number of regional factors included (the choice of n_f);
3. Whether the VAT data are included in the calculation of the regional factors or not, and the assumed publication lag of the VAT data;
4. How the regional factors are selected.

The first three differences listed above should be clear, but point four requires additional explanation. EMPCA, TW, and TP all involve using PCA. PCA using a data set of k_r variables will produce k_r factors. These are typically ordered (and selected for inclusion in the VAR) according to the proportion of the variability in the data which they explain (that is, based on the eigenvalues of the sample covariance matrix). A small number of factors are typically chosen that account for most of the variance in the data; for example, see McCracken and Ng (2021). One of our models uses this approach. That is, we simply add into the VAR the $n_f \ll k_r$ factors with the highest eigenvalues. However, it is possible that the factors with the highest eigenvalues may not be the ones that are the most useful for nowcasting regional GVA growth. Accordingly, we also experiment with selecting for inclusion in the MF-VAR, not those factors with the highest eigenvalue but those with the highest (in-sample, updated recursively) correlation with UK GVA growth. In the body of the paper, we report results for a selection of these MF-FAVARs with a focus on the issue of factor construction. A complete set of results using all our models is given in Appendix B. Overall, we find a high degree of robustness.

A final element to set out is the typical release calendar for the regional output data. We mimic this calendar in the design of our out-of-sample exercise. The current release schedule is presented in Figure 1. We choose to time the estimation of our model each quarter to coincide with the release of the UK growth rate for the previous quarter: this is approximately in the middle of the following quarter. From Figure 1, we can see that, given the release delays in producing estimates of regional output growth, there are three quantities of interest that can be produced using our model.

- “Backcasts”: An estimate of regional growth produced in quarter τ (say, 2020Q2) but which relates to quarter $\tau - 2$ (say, 2019Q4);
- “Estimates”: An estimate of regional growth produced in quarter τ (say, 2020Q2) but which relates to quarter $\tau - 1$ (say, 2020Q1);




Figure 1: Typical Release Schedule for National and Regional Output Data in the UK

- “Nowcasts”: An estimate of regional growth produced in quarter τ (say, 2020Q2) but which relates to quarter τ (say, 2020Q2).

We produce “nowcasts” and “estimates” for each region every time we run the model, but given the release schedule, we only produce “backcasts” for the English regions and Wales (because, continuing the example in 2020Q2, we already had estimates of Scottish and Northern Irish growth in 2019Q4, as these were released in March and April of 2020, respectively). Our “nowcasts” are estimates of regional growth in the quarter for which we do not yet have official data for the UK as a whole. When producing our “estimates” and “backcasts,” we already have the official estimate of UK growth for the corresponding quarter and these data are included in the model. In calculating the factors, we respect the release calendar for the predictors as set out in Table 1.

4.2 Nowcasting Results: Out-of-Sample Evidence

Having set out the timing convention for the production of our estimates, nowcasts, and backcasts we now present an evaluation of the performance of our models in estimating regional GVA growth. Our out-of-sample evaluation period is 2014Q2 through 2021Q4.⁹ We present the results from a number of different model specifications, as described earlier. These include results from the following three models:

1. An MF-VAR model (our benchmark model with no short data);

⁹Due to the lack of availability of real-time data vintages for our variables, our recursive out-of-sample analysis has to be “pseudo” real time.

2. An MF-FAVAR model;
3. An MF-FAVAR model, where the VAT data are not included, but all of the other regional (short) predictors are considered when estimating the factors.

In all cases, we evaluate the accuracy of the point and density nowcasts, estimates, and backcasts using the root mean square forecast error (RMSFE) and the continuous ranked probability score (CRPS), respectively. Lower values of each of these metrics indicate improved accuracy.

Table 2 presents the RMSFE metrics by region for each model, namely, the benchmark MF-VAR and the three factor-augmented MF-VAR models (MF-FAVAR), one for each of the three-factor estimation algorithms, produced using the adaptive Lasso-asymmetric conjugate prior. Several conclusions can be drawn from these results.

There is a clear pattern of accuracy improving as we move from our nowcast to our estimates to our backcasts. This reflects the accumulation of information that takes place between the production of each of these estimates over a given quarter. However, it is also notable that there is a much larger improvement in model performance as we go from the nowcast to the estimate, and a smaller improvement as we move from the estimate to the backcast. This is consistent with the finding in Koop et al. (Forthcoming), and reflects the fact that we know the aggregate (UK) estimate for that quarter when we produce our regional estimate (but not our regional nowcast). Crucially, it also reflects the presence of the additional measurement equation (which we refer to as the “cross-sectional restriction”) relating regional growth to aggregate national growth. Comparing the accuracy of the models including the additional short indicators set out earlier in this paper against the benchmark model, we generally see little or no improvement in the accuracy of the estimates or the backcasts. But there is an improvement in the accuracy of the nowcasts. Adding in the short data then helps. This conclusion holds across the different factor estimation algorithms (EMPCA, TW, and TP).

These conclusions hold when we evaluate the density nowcasts, estimates, and backcasts in Table 3 using the CRPS. Using Diebold and Mariano (1995) tests, we explore whether there are any statistically significant improvements in individual regions using the different MF-FAVAR models. For the density estimates and backcasts there are no statistically significant improvements, but the density nowcasts are almost always statistically significantly more accurate. This makes sense and is again consistent with the existing literature, in particular Koop et al. (Forthcoming). It reflects the fact that the largest improvement in the accuracy of our estimates and backcasts comes from conditioning directly on the equivalent UK estimate (which itself reflects much of the information contained in the additional predictors). When we make the prediction of regional growth earlier and, as a result, we do not yet know the UK outturn for a given quarter, the additional indicators, as captured by the factors, lead to substantial and statistically significant improvements in the accuracy of our nowcasts.

Table 2: RMSFE by Region (Multiplied by 100)

Benchmark MF-VAR														
	NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average	
Nowcast	2.84	1.82	2.10	1.87	1.96	2.08	3.35	1.80	1.72	2.46	1.63	1.71	2.11	
Estimate	0.84	0.48	0.54	0.58	0.41	0.38	1.21	0.40	0.57	0.68	0.16	0.38	0.55	
Backcast	0.42	0.30	0.35	0.36	0.26	0.27	0.79	0.19	0.31	0.34	-	-	0.36	
MF-FAVAR: model including VAT data														
EMPCA	Nowcast	1.68	1.62	1.67	1.72	1.61	1.64	2.45	1.52	1.54	1.82	1.24	1.36	1.66
	Estimate	0.62	0.55	0.62	0.67	0.45	0.40	1.10	0.41	0.59	0.62	0.15	0.33	0.54
	Backcast	0.33	0.31	0.38	0.41	0.27	0.26	0.70	0.21	0.35	0.32	-	-	0.35
TW	Nowcast	1.71	1.62	1.67	1.69	1.61	1.64	2.45	1.53	1.54	1.83	1.23	1.36	1.66
	Estimate	0.62	0.55	0.62	0.66	0.45	0.40	1.10	0.42	0.59	0.63	0.15	0.33	0.54
	Backcast	0.33	0.31	0.38	0.41	0.28	0.25	0.70	0.22	0.35	0.32	-	-	0.35
TP	Nowcast	1.72	1.63	1.66	1.70	1.61	1.65	2.44	1.53	1.54	1.82	1.23	1.38	1.66
	Estimate	0.62	0.56	0.62	0.67	0.46	0.41	1.10	0.41	0.59	0.63	0.15	0.34	0.55
	Backcast	0.33	0.31	0.38	0.41	0.28	0.26	0.70	0.21	0.35	0.32	-	-	0.36

Notes: NE, North East England; NW, North West England; York, Yorkshire and the Humber; EM, East Midlands; WM, West Midlands; EE, East of England; LON, London; SE, South East England; SW, South West England; WA, Wales; SCOT; Scotland; NI, Northern Ireland. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table 3: Average CRPS by Region (Multiplied by 100)

Benchmark MF-VAR													
	NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
Nowcast	1.21	0.87	0.99	0.91	0.89	0.91	1.23	0.84	0.86	1.07	0.61	0.72	0.93
Estimate	0.43	0.29	0.31	0.33	0.27	0.26	0.51	0.26	0.32	0.37	0.11	0.20	0.30
Backcast	0.23	0.17	0.19	0.19	0.16	0.15	0.28	0.14	0.18	0.20	-	-	0.19
MF-FAVAR: model including VAT data													
EMPCA	Nowcast	0.85*	0.81*	0.84	0.85	0.79*	1.05*	0.74*	0.79*	0.88*	0.51*	0.59*	0.79
	Estimate	0.36	0.31	0.33	0.35	0.29	0.49	0.26	0.33	0.36	0.11	0.19	0.30
	Backcast	0.20	0.18	0.20	0.21	0.17	0.27	0.14	0.19	0.19	-	-	0.19
TW	Nowcast	0.87*	0.81*	0.84	0.84	0.78*	1.05*	0.75*	0.79*	0.89*	0.50*	0.59*	0.79
	Estimate	0.36	0.31	0.33	0.35	0.29	0.49	0.26	0.33	0.36	0.11	0.19	0.30
	Backcast	0.20	0.18	0.20	0.21	0.17	0.27	0.14	0.19	0.19	-	-	0.19
TP	Nowcast	0.87*	0.82*	0.84	0.84	0.79*	1.04*	0.75*	0.79*	0.89*	0.50*	0.59*	0.79
	Estimate	0.36	0.31	0.33	0.35	0.29	0.49	0.26	0.33	0.36	0.11	0.19	0.30
	Backcast	0.20	0.18	0.20	0.21	0.17	0.27	0.14	0.19	0.19	-	-	0.19

Notes: NE, North East England; NW, North West England; York, Yorkshire and the Humber; EM, East Midlands; WM, West Midlands; EE, East of England; LON, London; SE, South East England; SW, South West England; WA, Wales; SCOT; Scotland; NI, Northern Ireland. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Having compared the performance of our FAVAR model to our benchmark MF-VAR, a specific question arises about the role of VAT data relative to other (short) regional indicators in improving the accuracy of the regional backcasts, estimates, and nowcasts. In order to explore this issue we reran the MF-FAVAR without the VAT data. The results from this additional exercise are presented in Tables 4 and 5. These show results that are consistent with those presented above; in particular, they bear the same result relative to the MF-VAR benchmark model, and in some cases, the RMSE/CRPS estimates are marginally better than those above. This is evidence that these VAT data are not adding significantly to our ability to nowcast regional GDP relative to our model with other regional predictors but no VAT data.

To check robustness to our modeling choices, in online Appendix B we present additional results covering cases where: (i) regional factors are not restricted to affect only their own region's output but are allowed to affect output in other regions too; (ii) we use a different prior (the adaptive Lasso); (iii) we use a different lag length in the VAR; (iv) we include a different number of factors in the model (5 rather than 3); (v) we select which factors to include based on their correlations with UK output growth rather than on the size of their eigenvalues; (vi) the regional VAT data are assumed to be available on a more timely basis than currently;¹⁰ and (vii) the cross-sectional restrictions are switched off. Apart from (vii), none of these modeling variants delivers consistent improvements in the accuracy of our different estimates relative to the results presented in the main paper. Indeed most do not markedly change the average evaluation metric across regions. The only version of these results where the evaluation metrics do change more substantially is when we switch off the cross-sectional restriction. In this case, the accuracy of our estimates (across different models) clearly gets worse.

¹⁰This is a counterfactual simulation exercise, designed to ascertain if the VAT data would be more useful if published more quickly than at present.

Table 4: RMSFE by Region (Multiplied by 100) when Excluding the VAT Data

MF-FAVAR, without VAT														
		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	Nowcast	1.72	1.59	1.66	1.69	1.61	1.63	2.24	1.57	1.54	1.87	1.23	1.40	1.65
	Estimate	0.58	0.53	0.60	0.64	0.45	0.40	0.97	0.42	0.56	0.66	0.15	0.35	0.53
	Backcast	0.31	0.30	0.36	0.39	0.27	0.25	0.63	0.23	0.33	0.33	-	-	0.34
TW	Nowcast	1.74	1.61	1.66	1.69	1.62	1.64	2.23	1.59	1.56	1.90	1.24	1.40	1.66
	Estimate	0.60	0.54	0.61	0.64	0.44	0.39	0.97	0.42	0.57	0.67	0.15	0.35	0.53
	Backcast	0.32	0.30	0.37	0.39	0.27	0.24	0.63	0.22	0.34	0.34	-	-	0.34
TP	Nowcast	1.72	1.59	1.64	1.68	1.61	1.62	2.22	1.59	1.54	1.89	1.23	1.39	1.64
	Estimate	0.58	0.53	0.60	0.63	0.45	0.39	0.97	0.43	0.57	0.67	0.15	0.34	0.53
	Backcast	0.30	0.30	0.36	0.39	0.27	0.25	0.63	0.22	0.33	0.34	-	-	0.34

Notes: NE, North East England; NW, North West England; York, Yorkshire and the Humber; EM, East Midlands; WM, West Midlands; EE, East of England; LON, London; SE, South East England; SW, South West England; WA, Wales; SCOT; Scotland; NI, Northern Ireland.
 * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table 5: Average CRPS by Region (multiplied by 100) when Excluding the VAT Data

MF-FAVAR, without VAT													
	NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	Nowcast	0.84*	0.77*	0.80*	0.80	0.75*	0.98*	0.73*	0.76*	0.87*	0.49*	0.57*	0.76
	Estimate	0.32	0.28	0.30	0.32	0.26	0.45*	0.23	0.29	0.34	0.10	0.18	0.27
	Backcast	0.18	0.16	0.19	0.19	0.15	0.25	0.13	0.17	0.18	-	-	0.17
TW	Nowcast	0.85*	0.78*	0.80*	0.80	0.75*	0.98*	0.74*	0.76*	0.87*	0.49*	0.57*	0.76
	Estimate	0.33	0.28	0.30	0.31	0.26	0.45*	0.23	0.29	0.35	0.10	0.18	0.27
	Backcast	0.18	0.16	0.19	0.19	0.15	0.25	0.13	0.17	0.18	-	-	0.17
TP	Nowcast	0.84*	0.77*	0.79*	0.79	0.75*	0.97*	0.74*	0.76*	0.87*	0.48*	0.57*	0.76
	Estimate	0.32	0.27	0.29	0.31	0.26	0.45*	0.24	0.29	0.35	0.10	0.18	0.27
	Backcast	0.17	0.16	0.19	0.19	0.15	0.25	0.13	0.17	0.18	-	-	0.17

Notes: NE, North East England; NW, North West England; York, Yorkshire and the Humber; EM, East Midlands; WM, West Midlands; EE, East of England; LON, London; SE, South East England; SW, South West England; WA, Wales; SCOT; Scotland; NI, Northern Ireland. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

5 Conclusions

The twin factors of increasing interest in regional economic issues and advances in the availability of data (including administrative data) have led to the creation of new regional economic predictors and the promise of being able to better track regional economic activity. In the UK, we have seen a significant increase in regional economic data over the past decade and, on the administrative data side, this now includes payroll employment information from the PAYE tax system and VAT turnover data. While these data may provide an indication of what is happening to regional output, they are not a direct estimate of it. However, many of these data series could be useful for nowcasting regional GVA (GDP). A major barrier to doing so in practice is the relatively short time series that characterize these data that makes their use in many nowcasting models problematic.

In this paper, we apply and test different methods for estimating large VAR models with mixed-frequency data when some of the indicators included in the VAR only have “short” historical coverage. We capture the information in the short data by constructing regional factors from the short data using a well-known (EMPCA) and two more recently developed (TP and TW) algorithms. These algorithms estimate common factors from data that have missing data at both the beginning and the end of the sample. We then add these regional factors into a regional MF-VAR, so as to exploit the putative information in the short data.

We find that the differences across each of the three factor extraction methods are small. It does not matter which algorithm is used to estimate factors from data sets characterized by a ragged edge at the beginning, as well as at the end, of the sample. When interest lies in producing “backcasts” or “estimates” of regional growth (using the nomenclature adopted in this paper), we find that there is little to be gained from incorporating these additional regional predictors into our model (relative to our relatively sophisticated benchmark model). When interest instead lies in producing (the more timely) “nowcasts” of regional output growth, before official data for UK GDP become available, we do find that there are gains to conditioning regional nowcasts on the factors, irrespective of which algorithm is used to estimate the factors. This suggests that, in this case, there is some utility in the short data. But as time passes and information on aggregate (UK-wide) economic activity within the quarter becomes available, the value of the short regional indicators declines.

References

- Bai, Jushan and Serena Ng (2006). “Confidence Intervals for Diffusion Index Forecasts and Inference for Factor-Augmented Regressions.” *Econometrica*, 74(4), pp. 1133–1150. doi:<https://doi.org/10.1111/j.1468-0262.2006.00696.x>.
- Bai, Jushan and Serena Ng (2021). “Matrix Completion, Counterfactuals, and Factor Analysis of Missing Data.” *Journal of the American Statistical Association*, 116(536), pp. 1746–1763. doi:[10.1080/01621459.2021.1967163](https://doi.org/10.1080/01621459.2021.1967163).
- Bañbura, Marta and Michele Modugno (2014). “Maximum likelihood estimation of factor models on datasets with arbitrary pattern of missing data.” *Journal of applied econometrics*, 29(1), pp. 133–160.
- Brave, Scott A., R. Andrew Butters, and Alejandro Justiniano (2019). “Forecasting Economic Activity with Mixed Frequency BVARs.” *International Journal of Forecasting*, 35(4), pp. 1692–1707. doi:[10.1016/j.ijforecast.2019.02.010](https://doi.org/10.1016/j.ijforecast.2019.02.010).
- Cahan, Ercument, Jushan Bai, and Serena Ng (2023). “Factor-Based Imputation of Missing Values and Covariances in Panel Data of Large Dimensions.” *Journal of Econometrics*, 233(1), pp. 113–131. doi:[10.1016/j.jeconom.2022.01.006](https://doi.org/10.1016/j.jeconom.2022.01.006).
- Chan, Joshua (2022). “Asymmetric Conjugate Priors for Large Bayesian VARs.” *Quantitative Economics*, 13(3), pp. 1145–1169. doi:[10.3982/QE1381](https://doi.org/10.3982/QE1381).
- Chan, Joshua C. C., Aubrey Poon, and Dan Zhu (2021). “Efficient Estimation of State-Space Mixed-Frequency VARs: A Precision-Based Approach.” doi:[10.48550/arXiv.2112.11315](https://doi.org/10.48550/arXiv.2112.11315).
- Diebold, Francis X and Robert S Mariano (1995). “Comparing Predictive Accuracy.” *Journal of Business & Economic Statistics*, 20(1), pp. 134–144. doi:[10.1080/07350015.1995.10524599](https://doi.org/10.1080/07350015.1995.10524599).
- Eraker, Bjorn, Ching Wai (Jeremy) Chiu, Andrew T. Foerster, Tae Bong Kim, and Hernan D. Seoane (2014). “Bayesian Mixed Frequency VARs.” *Journal of Financial Econometrics*, 13(3), pp. 698–721. doi:[10.1093/jjfinec/nbu027](https://doi.org/10.1093/jjfinec/nbu027).
- Gefang, Deborah, Gary Koop, and Aubrey Poon (2020). “Computationally Efficient Inference in Large Bayesian Mixed Frequency VARs.” *Economics Letters*, 191, pp. 109 – 120. doi:[10.1016/j.econlet.2020.109120](https://doi.org/10.1016/j.econlet.2020.109120).
- Koop, Gary, Stuart McIntyre, and James Mitchell (2020a). “UK Regional Nowcasting Using a Mixed Frequency Vector Auto-Regressive model with Entropic Tilting.” *Journal of the Royal Statistical Society: Series A (Statistics in Society)*, 183(1), pp. 91–119. doi:[10.1111/rssa.12491](https://doi.org/10.1111/rssa.12491).
- Koop, Gary, Stuart McIntyre, James Mitchell, and Aubrey Poon (2020b). “Reconciled Estimates and Nowcasts of Regional Output in the UK.” *National Institute Economic Review*, 253, pp. R44–R59. doi:[10.1017/nie.2020.29](https://doi.org/10.1017/nie.2020.29).

- Koop, Gary, Stuart McIntyre, James Mitchell, and Aubrey Poon (2020c). “Regional Output Growth in the United Kingdom: More Timely and Higher Frequency Estimates from 1970.” *Journal of Applied Econometrics*, 35(2), pp. 176–197. doi:10.1002/jae.2748.
- Koop, Gary, Stuart McIntyre, James Mitchell, and Aubrey Poon (Forthcoming). “Using Stochastic Hierarchical Aggregation Constraints to Nowcast Regional Economic Aggregates.” *International Journal of Forecasting*. doi:10.1016/j.ijforecast.2022.04.002.
- Labonne, Paul and Martin Weale (2020). “Temporal Disaggregation of Overlapping Noisy Quarterly Data: Estimation of Monthly Output from UK Value-Added Tax Data.” *Journal of the Royal Statistical Society, Series A*, 183(3), pp. 1211–1230. doi:10.1111/rssa.12568.
- Leamer, Edward E (2007). “Housing is the business cycle.” Working Paper 13428, National Bureau of Economic Research. doi:10.3386/w13428. URL <http://www.nber.org/papers/w13428>.
- McCracken, Michael W. and Serena Ng (2021). “FRED-QD: A Quarterly Database for Macroeconomic Research.” *Federal Reserve Bank of St Louis Review*, 103(1), pp. 1–44. doi:10.20955/r.103.1-44.
- McCracken, Michael W., Michael T. Owyang, and Tatevik Sekhposyan (2021). “Real-Time Forecasting and Scenario Analysis Using a Large Mixed-Frequency Bayesian VAR.” *International Journal of Central Banking*, 17(5), pp. 1–41. URL <https://www.ijcb.org/journal/ijcb21q5a8.htm>.
- Schorfheide, Frank and Dongho Song (2015). “Real-Time Forecasting With a Mixed-Frequency VAR.” *Journal of Business & Economic Statistics*, 33(3), pp. 366–380. doi:10.1080/07350015.2014.954707.
- Stock, J and M Watson (2002). “Macroeconomic Forecasting Using Diffusion Indexes.” *Journal of Business & Economic Statistics*, 20(2), pp. 147–162. doi:10.1198/073500102317351921.
- Stock, J and M Watson (2016). “Chapter 8 - Dynamic Factor Models, Factor-Augmented Vector Autoregressions, and Structural Vector Autoregressions in Macroeconomics.” In John B. Taylor and Harald Uhlig, editors, *Handbook of Macroeconomics*, volume 2, pp. 415–525. Elsevier. doi:10.1016/bs.hesmac.2016.04.002.
- Wallis, Kenneth F. (1986). “Forecasting with an Econometric Model: The ‘Ragged Edge’ Problem.” *Journal of Forecasting*, 5(1), pp. 1–13. doi:10.1002/for.3980050102.
- Zou, Hui (2006). “The Adaptive Lasso and Its Oracle Properties.” *Journal of the American Statistical Association*, 101(476), pp. 1418–1429. doi:10.1198/016214506000000735.

Appendices - for online publication only

A Monte Carlo Study Evaluating the Three Factor Extraction Algorithms With a Ragged Edge at the Beginning of the Sample

We simulate data from the approximate factor model also used in Bańbura and Modugno (2014) to generate data with different patterns of missing data:

$$\begin{aligned} y_t &= \chi_t + \epsilon_t = \Lambda_0 f_t + \dots + \Lambda_s f_{t-s} + \epsilon_t, \\ f_t &= A f_{t-1} + u_t, & u_t &\sim i.i.d. \mathcal{N}(0, I_r), \\ \epsilon_t &= D \epsilon_{t-1} + v_t, & v_t &\sim i.i.d. \mathcal{N}(0, \Phi) \end{aligned}$$

$t = 1, \dots, T$, where $\Lambda_{ij,k} \sim i.i.d. \mathcal{N}(0, 1)$, $i = 1, \dots, n, j = 1, \dots, r, k = 0, \dots, s$,

$$A_{ij} = \begin{cases} \rho, & i = j \\ 0, & i \neq j \end{cases}, D_{ij} = \begin{cases} \alpha, & i = j \\ 0, & i \neq j \end{cases}, \Phi_{i,j} = \tau^{|i-j|} (1 - \alpha^2) \sqrt{\gamma_i \gamma_j},$$

$$\gamma_i = \frac{\beta_i}{1 - \beta_i} \frac{1}{1 - \rho^2} \sum_{k=0}^s \sum_{j=1}^r \Lambda_{ij,k}^2, \beta_i \sim i.i.d. U([0.1, 0.9]).$$

Parameter τ controls the degree of cross-correlation of the idiosyncratic component. As described in Bańbura and Modugno (2014), $\tau = 0$ satisfies the assumption of a diagonal spectral density matrix of the idiosyncratic component required for an exact factor model; $\tau > 0$ violates this assumption, but satisfies the condition of weak cross-correlation (for an approximate factor model). Parameter α controls the degree of serial correlation of the idiosyncratic component. Parameter ρ controls the degree of persistence of factors. Parameter β_i controls the signal-to-noise ratio for variable i . Parameter s controls whether the relationship between the factors and the observables is “truly” dynamic $s > 0$ or not $s = 0$.

We then estimate factors on these simulated data using, in turn, the EMPCA, TW, and TP algorithms. We do so for simulated data that differ on the following dimensions:

(1): Time-series sample lengths, T : $T = 50$, $T = 100$, and $T = 200$;

(2): Cross-sectional sizes, n : $n = 12$, $n = 32$, $n = 52$, and $n = 102$;

(3): Values for τ : $\tau = 0$, $\tau = 0.3$, $\tau = 0.6$, and $\tau = 0.9$;

(4): Number of missing variables: Only the first two variables, i , are assumed to be complete.

The remaining $n - 2$ variables are generated with missing values.

(5): Fraction of data that are missing at the beginning of the sample: The first 0 percent, first 20 percent, first 40 percent, first 60 percent, or first 80 percent of data are assumed to be missing.

For other parameters, we set them as fixed across different experiments: The true number of factors $r = 2$, the degree of persistence of factors $\rho = 0.7$, the degree of serial correlation $\alpha = 0.5$, the relationship between the factors, and the observables $s = 0$. Finally, in the estimation procedure, we set the estimated number of factors as $\hat{r} = r$.

To assess the precision of the estimates, as in Bańbura and Modugno (2014), we use the trace R^2

of the regression of the estimated factors (from EMPCA, TW, or TP) on the true ones (as simulated):

$$\frac{\text{Trace} \left(F' \hat{F} \left(\hat{F}' \hat{F} \right)^{-1} \hat{F}' F \right)}{\text{Trace} (F' F)},$$

where $\hat{F} = \mathbf{E}_{\hat{\theta}} [F \mid \Omega_T]$.

Table A1 shows the trace R^2 and the computation time using the EMPCA algorithm. “Not,” as reported in the table, means that, across the 500 replications, there is at least one replication where EMPCA does not converge after 5,000 iterations. The convergence criteria is the maximum percentage change in the variables’ estimates is smaller than 0.001. Table A2 provides analogous results for TW and TP.

As for the trace R^2 for the EMPCA, the estimates are less precise for small sample lengths ($T = 50$ versus $T = 200$), small cross-sections ($n = 12$ versus $n = 102$), a mis-specified model ($\tau = 0.9$ versus $\tau = 0$) in small samples, and a large fraction of missing data.

As for the computation time for the EMPCA, it takes longer for large sample lengths ($T = 200$ versus $T = 50$), large cross-sections ($n = 102$ versus $n = 12$), a mis-specified model ($\tau = 0.9$ versus $\tau = 0$) in small samples, and a large fraction of missing data. But the additional computational burden is small. However, there were some instances of convergence failures, especially for a mis-specified model ($\tau = 0.9$ versus $\tau = 0$) and a large fraction of missing data.

When we compare the trace R^2 across the EMPCA, TW, and TP algorithms, we find that the three algorithms all have similar estimation accuracy.

Table A1: EMPCA: Trace R^2 and Computation Times (in Seconds), where $r = 2$

T	n	τ	Trace R^2 EMPCA					Computation Time			
			missing: 0	20%	40%	60%	80%	20%	40%	60%	80%
50	12	0	0.77	0.59	0.53	0.47	0.40	0.19	0.30	0.23	0.29
50	12	0.3	0.66	0.49	0.46	0.41	0.38	0.23	0.35	0.27	0.30
50	12	0.6	0.51	0.37	0.34	0.32	0.35	0.29	0.41	0.27	0.32
50	12	0.9	0.49	0.32	0.31	0.34	0.34	0.27	0.34	0.29	0.34
50	32	0	0.89	0.66	0.54	0.49	0.41	1.27	1.68	1.66	1.84
50	32	0.3	0.85	0.61	0.49	0.41	0.40	1.36	1.87	1.83	1.88
50	32	0.6	0.75	0.48	0.43	0.40	0.34	1.67	2.23	2.04	1.91
50	32	0.9	0.43	0.32	0.32	0.32	0.31	2.30	2.57	2.16	1.98
50	52	0	0.91	0.64	0.56	0.47	0.44	2.69	3.63	3.26	3.11
50	52	0.3	0.89	0.61	0.53	0.43	0.38	3.35	3.91	3.38	3.27
50	52	0.6	0.84	0.53	0.45	0.39	0.37	4.03	4.26	3.62	3.37
50	52	0.9	0.48	0.35	0.28	0.32	0.37	4.89	5.14	3.97	not
50	102	0	0.94	0.64	0.59	0.46	0.40	9.93	9.70	7.86	not
50	102	0.3	0.92	0.64	0.54	0.44	0.41	10.03	10.52	8.20	not
50	102	0.6	0.89	0.59	0.47	0.42	0.37	10.81	11.10	8.47	not
50	102	0.9	0.63	0.36	0.37	0.36	0.36	13.69	12.49	not	not
100	12	0	0.82	0.63	0.55	0.43	0.43	0.70	0.86	1.03	0.83
100	12	0.3	0.72	0.55	0.46	0.44	0.38	0.78	1.13	1.18	0.87
100	12	0.6	0.48	0.35	0.36	0.34	0.33	1.14	1.26	1.23	0.93
100	12	0.9	0.50	0.33	0.35	0.32	0.38	0.98	1.12	1.14	0.84
100	32	0	0.92	0.65	0.57	0.50	0.45	4.59	4.89	4.71	3.69
100	32	0.3	0.88	0.66	0.54	0.44	0.42	5.20	5.99	5.39	3.72
100	32	0.6	0.79	0.55	0.46	0.41	0.38	6.10	6.64	5.97	4.01
100	32	0.9	0.42	0.30	0.28	0.34	0.35	7.94	7.50	6.32	not
100	52	0	0.94	0.70	0.58	0.50	0.42	9.79	10.80	not	not
100	52	0.3	0.92	0.65	0.52	0.44	0.41	11.40	11.57	not	not
100	52	0.6	0.88	0.59	0.53	0.38	0.38	18.37	not	not	not
100	52	0.9	0.44	0.30	0.31	0.32	0.35	17.42	not	not	not
100	102	0	0.96	0.68	0.60	0.47	0.45	29.43	not	not	not
100	102	0.3	0.95	0.65	0.57	0.46	0.43	31.30	not	not	not
100	102	0.6	0.93	0.66	0.52	0.43	0.37	31.87	not	not	not
100	102	0.9	0.68	0.42	0.37	0.32	0.33	not	not	not	not
200	12	0	0.84	0.63	0.55	0.50	0.45	3.35	3.82	3.79	2.85
200	12	0.3	0.76	0.57	0.45	0.43	0.40	4.53	5.57	4.79	3.04
200	12	0.6	0.48	0.36	0.37	0.34	0.34	6.02	6.37	5.38	3.56
200	12	0.9	0.48	0.32	0.34	0.32	0.35	5.45	5.42	4.65	3.23
200	32	0	0.93	0.72	0.57	0.53	0.45	22.23	22.99	17.19	11.85
200	32	0.3	0.90	0.67	0.55	0.45	0.42	23.98	23.21	19.09	not
200	32	0.6	0.83	0.55	0.47	0.40	0.35	34.82	28.91	22.93	not
200	32	0.9	0.42	0.30	0.30	0.32	0.29	41.70	35.42	not	not
200	52	0	0.96	0.78	0.60	0.63	0.52	28.64	39.64	40.00	not
200	52	0.3	0.94	0.53	0.56	0.42	0.46	61.17	29.83	38.05	not
200	52	0.6	0.91	0.50	0.46	0.34	0.35	83.58	37.23	not	not
200	52	0.9	0.51	0.23	0.22	0.36	0.40	83.58	not	not	not
200	102	0	0.97	0.84	0.59	0.71	0.55	114.13	not	not	not
200	102	0.3	0.97	0.58	0.65	0.28	0.48	150.03	not	not	not
200	102	0.6	0.95	0.61	0.49	0.46	0.50	161.98	not	not	not
200	102	0.9	0.78	0.46	0.35	0.33	0.24	not	not	not	not

Table A2: Trace R^2 : TW and TP, where $r = 2$

T	n	τ	Trace R^2 TW					Trace R^2 TP				
			missing: 0	20%	40%	60%	80%	missing: 0	20%	40%	60%	80%
50	12	0	0.79	0.69	0.61	0.48	0.40	0.79	0.71	0.59	0.49	0.44
50	12	0.3	0.68	0.61	0.52	0.42	0.38	0.68	0.62	0.56	0.48	0.42
50	12	0.6	0.52	0.44	0.41	0.35	0.31	0.48	0.50	0.42	0.38	0.35
50	12	0.9	0.47	0.45	0.39	0.35	0.29	0.51	0.47	0.41	0.37	0.38
50	32	0	0.89	0.78	0.66	0.54	0.40	0.88	0.79	0.68	0.53	0.44
50	32	0.3	0.86	0.75	0.62	0.49	0.38	0.84	0.74	0.63	0.51	0.40
50	32	0.6	0.73	0.63	0.54	0.43	0.34	0.73	0.64	0.57	0.44	0.39
50	32	0.9	0.42	0.37	0.35	0.29	0.27	0.38	0.38	0.37	0.35	0.38
50	52	0	0.91	0.79	0.67	0.53	0.40	0.91	0.78	0.67	0.55	0.44
50	52	0.3	0.89	0.78	0.66	0.50	0.39	0.90	0.78	0.67	0.53	0.41
50	52	0.6	0.84	0.72	0.57	0.45	0.31	0.83	0.71	0.60	0.48	0.38
50	52	0.9	0.49	0.42	0.39	0.35	0.27	0.46	0.42	0.41	0.39	0.35
50	102	0	0.93	0.81	0.69	0.54	0.38	0.92	0.81	0.70	0.56	0.43
50	102	0.3	0.91	0.80	0.67	0.53	0.38	0.92	0.80	0.69	0.53	0.43
50	102	0.6	0.90	0.76	0.64	0.46	0.36	0.89	0.76	0.64	0.48	0.39
50	102	0.9	0.65	0.58	0.47	0.41	0.27	0.65	0.57	0.50	0.42	0.38
100	12	0	0.83	0.72	0.63	0.53	0.43	0.83	0.74	0.63	0.54	0.47
100	12	0.3	0.73	0.64	0.56	0.47	0.39	0.73	0.66	0.59	0.47	0.43
100	12	0.6	0.51	0.47	0.42	0.35	0.33	0.49	0.48	0.41	0.39	0.40
100	12	0.9	0.53	0.47	0.39	0.35	0.34	0.44	0.44	0.44	0.37	0.32
100	32	0	0.92	0.81	0.70	0.57	0.46	0.92	0.82	0.69	0.61	0.46
100	32	0.3	0.89	0.79	0.65	0.54	0.40	0.89	0.78	0.66	0.56	0.42
100	32	0.6	0.80	0.70	0.58	0.44	0.35	0.80	0.71	0.60	0.48	0.39
100	32	0.9	0.43	0.38	0.34	0.30	0.28	0.36	0.39	0.38	0.35	0.38
100	52	0	0.94	0.82	0.71	0.60	0.47	0.93	0.83	0.72	0.62	0.48
100	52	0.3	0.92	0.81	0.69	0.56	0.40	0.93	0.82	0.70	0.56	0.42
100	52	0.6	0.89	0.76	0.64	0.51	0.35	0.88	0.77	0.66	0.52	0.40
100	52	0.9	0.47	0.41	0.38	0.33	0.29	0.44	0.49	0.43	0.44	0.35
100	102	0	0.96	0.84	0.72	0.58	0.46	0.96	0.85	0.73	0.62	0.48
100	102	0.3	0.95	0.83	0.70	0.55	0.42	0.95	0.83	0.71	0.58	0.46
100	102	0.6	0.93	0.80	0.66	0.54	0.38	0.93	0.81	0.67	0.53	0.39
100	102	0.9	0.71	0.62	0.51	0.40	0.30	0.70	0.63	0.59	0.45	0.37
200	12	0	0.84	0.74	0.65	0.54	0.47	0.85	0.75	0.65	0.56	0.43
200	12	0.3	0.76	0.68	0.57	0.48	0.40	0.76	0.68	0.58	0.52	0.44
200	12	0.6	0.55	0.45	0.41	0.35	0.35	0.54	0.47	0.46	0.40	0.38
200	12	0.9	0.50	0.44	0.39	0.35	0.32	0.49	0.44	0.41	0.40	0.37
200	32	0	0.93	0.83	0.72	0.59	0.48	0.94	0.83	0.72	0.62	0.51
200	32	0.3	0.90	0.80	0.67	0.57	0.42	0.91	0.81	0.70	0.58	0.45
200	32	0.6	0.84	0.75	0.60	0.49	0.38	0.85	0.73	0.63	0.51	0.40
200	32	0.9	0.37	0.37	0.30	0.30	0.26	0.42	0.36	0.35	0.36	0.39
200	52	0	0.96	0.85	0.72	0.62	0.48	0.95	0.84	0.73	0.62	0.49
200	52	0.3	0.94	0.82	0.70	0.58	0.47	0.94	0.83	0.71	0.60	0.47
200	52	0.6	0.91	0.79	0.67	0.53	0.40	0.90	0.80	0.66	0.53	0.41
200	52	0.9	0.40	0.35	0.35	0.30	0.24	0.40	0.40	0.41	0.43	0.38
200	102	0	0.97	0.87	0.74	0.61	0.49	0.97	0.86	0.74	0.64	0.51
200	102	0.3	0.96	0.85	0.73	0.59	0.48	0.96	0.85	0.73	0.61	0.49
200	102	0.6	0.95	0.84	0.69	0.57	0.44	0.95	0.83	0.70	0.57	0.43
200	102	0.9	0.76	0.66	0.55	0.41	0.31	0.73	0.69	0.56	0.48	0.40

B Nowcasting Results: Robustness Checks

In this appendix, as summarized in the main paper, we provide additional nowcasting results that provide robustness checks on our main model specification (used in the main paper). Key dimensions to which we assess the robustness are: 1) Different inclusions of impacts (every region impacting on all others; 2) different priors (adaptive Lasso prior); 3) different VAR lag lengths ($p=4$ lags); 4) a different number of factors ($n_f = 5$); 5) different ways of selecting the factors (according to the correlation with UK GVA); 6) simulating a reduction in the release delay for the VAT data; and 7) if the cross-sectional restrictions are switched off.

Results for these 7 robustness checks are presented, in turn, in the following 7 sections.

1 Allowing Every Region to Impact All Others

Table B1: RMSFE (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-full - AL-ACP (p=1 lag, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	1.69	1.60	1.66	1.70	1.59	1.60	2.45	1.50	1.52	1.77	1.21	1.29	1.63
		Estimate	0.64	0.56	0.63	0.68	0.46	0.40	1.12	0.43	0.62	0.61	0.15	0.32	0.55
		Backcast	0.34	0.31	0.38	0.42	0.28	0.26	0.71	0.23	0.35	0.31	-	-	0.36
TW	AL-ACP	Nowcast	1.70	1.59	1.65	1.68	1.60	1.60	2.42	1.50	1.51	1.78	1.20	1.30	1.63
		Estimate	0.63	0.56	0.63	0.67	0.46	0.40	1.11	0.42	0.60	0.62	0.15	0.33	0.55
		Backcast	0.34	0.31	0.38	0.41	0.28	0.25	0.70	0.22	0.34	0.32	-	-	0.36
TP	AL-ACP	Nowcast	1.71	1.59	1.65	1.67	1.60	1.61	2.41	1.49	1.50	1.78	1.20	1.30	1.63
		Estimate	0.63	0.56	0.62	0.67	0.47	0.40	1.10	0.41	0.60	0.62	0.15	0.33	0.55
		Backcast	0.34	0.31	0.38	0.41	0.28	0.26	0.70	0.22	0.35	0.32	-	-	0.36

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table B2: Average CRPS (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-full - AL-ACP (p=1 lag, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	0.86*	0.81*	0.83	0.84	0.78*	0.79*	1.06*	0.73*	0.78*	0.87*	0.50*	0.57*	0.79
		Estimate	0.36	0.31	0.34	0.36	0.29	0.26	0.50	0.27	0.34	0.35	0.11	0.19	0.31
		Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.28	0.15	0.19	0.19	-	-	0.19
TW	AL-ACP	Nowcast	0.86*	0.81*	0.83	0.83	0.78*	0.79*	1.05*	0.74*	0.78*	0.88*	0.50*	0.58*	0.78
		Estimate	0.36	0.31	0.33	0.35	0.29	0.26	0.49	0.27	0.33	0.36	0.11	0.19	0.30
		Backcast	0.20	0.18	0.21	0.21	0.17	0.15	0.27	0.15	0.19	0.19	-	-	0.19
TP	AL-ACP	Nowcast	0.87*	0.81*	0.83	0.83	0.78*	0.79*	1.05*	0.73*	0.78*	0.88*	0.50*	0.58*	0.78
		Estimate	0.36	0.31	0.33	0.35	0.30	0.26	0.50	0.26	0.33	0.36	0.11	0.19	0.31
		Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

2 A Different Prior: Adaptive Lasso (AL) Prior

Table B3: RMSFE (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-own - AL (p=1 lag, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL	Nowcast	1.74	1.63	1.71	1.70	1.69	1.88	2.53	1.80	1.60	1.60	1.20	1.42	1.71
		Estimate	0.81	0.52	0.53	0.57	0.43	0.49	1.14	0.49	0.55	0.54	0.16	0.35	0.55
		Backcast	0.46	0.28	0.31	0.35	0.27	0.27	0.72	0.24	0.32	0.28	-	-	0.35
TW	AL	Nowcast	1.74	1.60	1.70	1.67	1.69	1.86	2.51	1.77	1.60	1.58	1.20	1.40	1.69
		Estimate	0.81	0.51	0.53	0.56	0.44	0.50	1.13	0.49	0.55	0.54	0.16	0.35	0.55
		Backcast	0.45	0.27	0.31	0.34	0.27	0.27	0.72	0.24	0.32	0.28	-	-	0.35
TP	AL	Nowcast	1.74	1.61	1.65	1.65	1.67	1.86	2.53	1.79	1.57	1.60	1.20	1.39	1.69
		Estimate	0.80	0.52	0.54	0.58	0.43	0.50	1.15	0.49	0.56	0.54	0.16	0.34	0.55
		Backcast	0.45	0.27	0.32	0.36	0.27	0.27	0.72	0.24	0.33	0.28	-	-	0.35

Notes: * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table B4: Average CRPS (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-own - AL (p=1 lag, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL	Nowcast	0.84	0.81*	0.86*	0.88	0.83*	0.81*	1.10*	0.87*	0.88	0.86*	0.52	0.62*	0.82
		Estimate	0.41	0.30	0.31*	0.35*	0.29	0.25	0.51	0.30*	0.36	0.35	0.11	0.19	0.31
		Backcast	0.24	0.17	0.18	0.21*	0.17	0.14	0.28	0.16	0.21*	0.19	-	-	0.19
TW	AL	Nowcast	0.85	0.80*	0.85*	0.87	0.83*	0.81*	1.10*	0.86*	0.88	0.85*	0.52	0.62*	0.31
		Estimate	0.41	0.29	0.32	0.35	0.29	0.25	0.51	0.30*	0.35	0.35	0.11	0.20	0.19
		Backcast	0.24	0.16	0.18	0.21	0.17	0.14	0.28	0.16	0.21	0.19	-	-	0.19
TP	AL	Nowcast	0.85	0.80*	0.84*	0.86	0.82*	0.81*	1.10*	0.86*	0.87	0.86*	0.52	0.61*	0.82
		Estimate	0.41	0.30	0.32*	0.35*	0.29	0.25	0.51	0.30*	0.36	0.35	0.11	0.19	0.31
		Backcast	0.24	0.16	0.19	0.21*	0.17	0.14	0.28	0.16	0.21*	0.19	-	-	0.19

Notes: * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

3 A Different VAR Lag Length: $p=4$ Lags

Table B5: RMSFE (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-own - AL-ACP ($p=4$ lags, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	1.87	1.69	1.86	1.98	1.71	1.76	2.67	1.69	1.68	1.88	1.35	1.43	1.80
		Estimate	0.74	0.58	0.84	1.06	0.59	0.41	1.18	0.44	0.73	0.63	0.16	0.34	0.64
		Backcast	0.40	0.33	0.50	0.66	0.36	0.27	0.75	0.21	0.44	0.32	-	-	0.42
TW	AL-ACP	Nowcast	1.88	1.70	1.85	1.98	1.70	1.76	2.66	1.69	1.69	1.88	1.35	1.43	1.80
		Estimate	0.75	0.59	0.84	1.06	0.58	0.41	1.18	0.44	0.73	0.63	0.16	0.34	0.64
		Backcast	0.40	0.33	0.51	0.66	0.36	0.27	0.75	0.21	0.44	0.32	-	-	0.43
TP	AL-ACP	Nowcast	1.87	1.70	1.85	1.99	1.71	1.76	2.66	1.70	1.69	1.88	1.35	1.43	1.80
		Estimate	0.74	0.59	0.84	1.05	0.59	0.41	1.17	0.44	0.73	0.63	0.16	0.34	0.64
		Backcast	0.40	0.33	0.50	0.66	0.36	0.27	0.74	0.21	0.44	0.32	-	-	0.42

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table B6: Average CRPS (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-own - AL-ACP ($p=4$ lags, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	0.94	0.86	0.95	0.99	0.86	0.87	1.21	0.85	0.86	0.92*	0.59	0.64	0.88
		Estimate	0.41	0.32	0.43*	0.52*	0.34	0.27	0.54	0.27	0.38*	0.36	0.11	0.19	0.35
		Backcast	0.23	0.19	0.26	0.32	0.20	0.16	0.31	0.15	0.23	0.19	-	-	0.22
TW	AL-ACP	Nowcast	0.94	0.86	0.94	0.99	0.85	0.87	1.20	0.85	0.86	0.92*	0.59	0.64	0.88
		Estimate	0.41	0.32	0.43*	0.52*	0.34	0.27	0.54	0.27	0.38*	0.36	0.11	0.19	0.34
		Backcast	0.23	0.18	0.26	0.32	0.20	0.16	0.31	0.15	0.23	0.19	-	-	0.22
TP	AL-ACP	Nowcast	0.94	0.86	0.94	0.99	0.85	0.87	1.20	0.85	0.86	0.92*	0.59	0.64	0.88
		Estimate	0.40	0.32	0.43*	0.52*	0.34	0.27	0.54	0.27	0.38*	0.36	0.11	0.19	0.34
		Backcast	0.23	0.19	0.26	0.32	0.20	0.16	0.30	0.14	0.23	0.19	-	-	0.22

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

4 A Different Number of Factors: $n_f = 5$ factors

Table B7: RMSFE (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-own - AL-ACP (p=1 lag, $n_f=5$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	1.69	1.61	1.67	1.71	1.60	1.63	2.46	1.52	1.53	1.82	1.24	1.36	1.65
		Estimate	0.62	0.55	0.63	0.67	0.45	0.40	1.10	0.41	0.59	0.62	0.15	0.33	0.55
		Backcast	0.34	0.31	0.38	0.42	0.28	0.25	0.70	0.22	0.35	0.32	-	-	0.36
TW	AL-ACP	Nowcast	1.71	1.62	1.68	1.71	1.61	1.64	2.45	1.53	1.54	1.82	1.23	1.37	1.66
		Estimate	0.61	0.56	0.63	0.67	0.45	0.40	1.11	0.41	0.59	0.62	0.15	0.34	0.55
		Backcast	0.33	0.31	0.38	0.42	0.28	0.26	0.70	0.22	0.35	0.32	-	-	0.36
TP	AL-ACP	Nowcast	1.71	1.61	1.66	1.70	1.60	1.63	2.44	1.54	1.53	1.82	1.23	1.36	1.65
		Estimate	0.62	0.56	0.62	0.67	0.46	0.40	1.10	0.42	0.60	0.63	0.15	0.34	0.55
		Backcast	0.34	0.31	0.38	0.41	0.28	0.25	0.70	0.22	0.35	0.32	-	-	0.36

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table B8: Average CRPS (Multiplied by 100), with Factors Selected Based on the Size of the Eigenvalue

		VB MF-FAVAR-own - AL-ACP (p=1 lag, $n_f=5$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	0.86*	0.81*	0.84	0.84	0.78*	0.79*	1.06*	0.74*	0.79*	0.88*	0.50*	0.59*	0.79
		Estimate	0.36	0.31	0.34	0.35	0.29	0.26	0.49	0.26	0.33	0.36	0.11	0.19	0.30
		Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19
TW	AL-ACP	Nowcast	0.86*	0.81*	0.84	0.84	0.79*	0.79*	1.05*	0.75*	0.79*	0.89*	0.50*	0.59*	0.79
		Estimate	0.35	0.31	0.33	0.35	0.29	0.26	0.50	0.26	0.33	0.36	0.11	0.19	0.30
		Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19
TP	AL-ACP	Nowcast	0.87*	0.81*	0.83*	0.84	0.78*	0.79*	1.05*	0.74*	0.79*	0.89*	0.50*	0.59*	0.79
		Estimate	0.36	0.31	0.33	0.35*	0.29	0.26	0.50	0.26	0.33	0.36	0.11	0.19	0.30
		Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.28	0.14	0.19	0.19	-	-	0.19

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

5 Selecting the Factors Based on their Correlation with UK GDP Growth

Table B9: RMSFE (Multiplied by 100)

		VB MF-FAVAR-own - AL-ACP (p=1 lag, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	1.70	1.63	1.68	1.73	1.61	1.64	2.45	1.53	1.55	1.83	1.24	1.36	1.66
		Estimate	0.62	0.55	0.63	0.68	0.46	0.40	1.10	0.41	0.60	0.62	0.15	0.34	0.55
		Backcast	0.33	0.31	0.38	0.42	0.28	0.25	0.70	0.22	0.35	0.31	-	-	0.36
TW	AL-ACP	Nowcast	1.71	1.63	1.67	1.70	1.62	1.63	2.44	1.53	1.54	1.83	1.24	1.37	1.66
		Estimate	0.62	0.56	0.63	0.67	0.46	0.40	1.10	0.41	0.59	0.62	0.15	0.34	0.55
		Backcast	0.33	0.31	0.38	0.41	0.28	0.25	0.70	0.22	0.35	0.32	-	-	0.36
TP	AL-ACP	Nowcast	1.72	1.63	1.67	1.70	1.61	1.64	2.43	1.54	1.53	1.84	1.23	1.36	1.66
		Estimate	0.62	0.55	0.62	0.67	0.45	0.40	1.09	0.41	0.59	0.64	0.15	0.34	0.55
		Backcast	0.33	0.31	0.38	0.41	0.27	0.25	0.69	0.22	0.35	0.32	-	-	0.35

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table B10: Average CRPS (Multiplied by 100)

		VB MF-FAVAR-own - AL-ACP (p=1 lag, $n_f=2$, VAT lag = 5 months)													
F_est	Prior		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
EMPCA	AL-ACP	Nowcast	0.87*	0.81*	0.84	0.85	0.79*	0.79*	1.05*	0.75*	0.80*	0.88*	0.50*	0.59*	0.79
		Estimate	0.36	0.31	0.34	0.35	0.29	0.26	0.49	0.26	0.33	0.35	0.11	0.19	0.30
		Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19
TW	AL-ACP	Nowcast	0.87*	0.82*	0.84	0.84	0.79*	0.79*	1.05*	0.74*	0.80*	0.88*	0.50*	0.59*	0.79
		Estimate	0.36	0.31	0.34	0.35	0.29	0.26	0.49	0.26	0.33	0.36	0.11	0.19	0.30
		Backcast	0.20	0.18	0.21	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19
TP	AL-ACP	Nowcast	0.87*	0.81*	0.83	0.84	0.79*	0.79*	1.04*	0.74*	0.79*	0.89*	0.50*	0.59*	0.79
		Estimate	0.36	0.31	0.33	0.35	0.29	0.26	0.49	0.26	0.33	0.36	0.11	0.19	0.30
		Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19

Notes: AL-ACP means Adaptive Lasso - Asymmetric Conjugate Prior. * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

6 Simulating a Reduction in the Release Delay for the Regional VAT Data

Table B11: RMSFE (Multiplied by 100)

		Benchmark MF-VAR												
		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
	Nowcast	2.84	1.82	2.10	1.87	1.96	2.08	3.35	1.80	1.72	2.46	1.63	1.71	2.11
	Estimate	0.84	0.48	0.54	0.58	0.41	0.38	1.21	0.40	0.57	0.68	0.16	0.38	0.55
	Backcast	0.42	0.30	0.35	0.36	0.26	0.27	0.79	0.19	0.31	0.34	-	-	0.36
		MF-FAVAR: w/ VAT and lag=1 month												
EMPCA	Nowcast	1.70	1.62	1.67	1.71	1.61	1.63	2.45	1.53	1.54	1.82	1.24	1.37	1.66
	Estimate	0.62	0.55	0.62	0.67	0.45	0.41	1.10	0.41	0.59	0.62	0.15	0.34	0.55
	Backcast	0.33	0.31	0.38	0.42	0.27	0.26	0.70	0.22	0.35	0.31	-	-	0.35
TW	Nowcast	1.72	1.64	1.66	1.72	1.61	1.64	2.44	1.54	1.53	1.82	1.23	1.36	1.66
	Estimate	0.63	0.55	0.62	0.68	0.46	0.41	1.10	0.42	0.59	0.62	0.15	0.34	0.55
	Backcast	0.33	0.31	0.38	0.42	0.28	0.26	0.70	0.22	0.35	0.32	-	-	0.35
TP	Nowcast	1.71	1.62	1.65	1.70	1.61	1.63	2.44	1.54	1.54	1.83	1.23	1.36	1.66
	Estimate	0.63	0.55	0.62	0.66	0.46	0.40	1.10	0.42	0.59	0.63	0.15	0.33	0.55
	Backcast	0.34	0.31	0.38	0.41	0.28	0.25	0.70	0.22	0.35	0.32	-	-	0.36

Notes: * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

Table B12: Average CRPS (Multiplied by 100)

		Benchmark MF-VAR												
		NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
	Nowcast	1.21	0.87	0.99	0.91	0.89	0.91	1.23	0.84	0.86	1.07	0.61	0.72	0.93
	Estimate	0.43	0.29	0.31	0.33	0.27	0.26	0.51	0.26	0.32	0.37	0.11	0.20	0.30
	Backcast	0.23	0.17	0.19	0.19	0.16	0.15	0.28	0.14	0.18	0.20	-	-	0.19
		MF-FAVAR: w/ VAT and lag=1 month												
EMPCA	Nowcast	0.86*	0.81*	0.84	0.84	0.79*	0.79*	1.05*	0.74*	0.79*	0.88*	0.50*	0.60*	0.79
	Estimate	0.36	0.31	0.33	0.36	0.29	0.26	0.49	0.26	0.33	0.36	0.11	0.19	0.30
	Backcast	0.20	0.17	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19
TW	Nowcast	0.87*	0.82*	0.84	0.85	0.79*	0.80*	1.05*	0.75*	0.79*	0.89*	0.50*	0.59*	0.79
	Estimate	0.36	0.31	0.33	0.35	0.29	0.27	0.49	0.26	0.33	0.36	0.11	0.19	0.30
	Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19
TP	Nowcast	0.87*	0.81*	0.83*	0.84	0.79*	0.79*	1.05*	0.74*	0.79*	0.89*	0.50*	0.59*	0.79
	Estimate	0.36	0.31	0.33	0.35	0.29	0.26	0.49	0.26	0.33	0.36	0.11	0.19	0.30
	Backcast	0.20	0.18	0.20	0.21	0.17	0.15	0.27	0.14	0.19	0.19	-	-	0.19

Notes: * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test.

7 Switching Off the Cross-Sectional Restrictions

Table B13: RMSFE (Multiplied by 100)

Benchmark MF-VAR without cross-sectional restrictions													
	NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
Nowcast	2.52	2.15	2.33	2.12	2.28	2.14	1.81	2.17	2.11	1.84	1.97	1.72	2.10
Estimate	1.49	1.45	1.47	1.36	1.41	1.34	1.11	1.41	1.38	1.09	0.69	0.79	1.25
Backcast	0.84	0.89	0.88	0.83	0.81	0.85	0.61	0.83	0.80	0.66	-	-	0.67
MF-FAVAR: no VAT data													
Nowcast	2.25	1.93	2.25	2.13	2.17	1.95	1.67	2.01	2.07	1.77	1.95	1.73	1.99
Estimate	1.28	1.30	1.39	1.35	1.33	1.20	1.00	1.27	1.32	0.97	0.68	0.76	1.15
Backcast	0.71	0.79	0.81	0.80	0.74	0.75	0.53	0.73	0.74	0.55	-	-	0.60
MF-FAVAR: with VAT data													
Nowcast	2.17	2.02	2.22	2.28	2.41	2.15	1.62	2.15	2.15	1.58	1.62	1.42	1.98
Estimate	1.27	1.31	1.40	1.43	1.45	1.33	0.97	1.36	1.36	0.85	0.69	0.66	1.17
Backcast	0.69	0.81	0.83	0.85	0.81	0.81	0.52	0.81	0.80	0.45	-	-	0.62
MF-FAVAR: with VAT data and lag=1 month													
Nowcast	2.17	2.02	2.23	2.29	2.41	2.15	1.62	2.14	2.14	1.59	1.62	1.43	1.98
Estimate	1.27	1.31	1.40	1.43	1.46	1.33	0.97	1.36	1.36	0.84	0.69	0.67	1.17
Backcast	0.69	0.81	0.83	0.86	0.81	0.81	0.52	0.81	0.81	0.45	-	-	0.62

Notes: * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test. All models do not have the cross-sectional restrictions imposed. All factors are estimated using the TP approach.

Table B14: Average CRPS (Multiplied by 100)

Benchmark MF-VAR without cross-sectional restrictions													
	NE	NW	York	EM	WM	EE	LON	SE	SW	WA	SCOT	NI	Average
Nowcast	1.21	1.07	1.10	1.01	1.08	1.03	0.93	1.05	0.99	0.94	0.74	0.73	0.99
Estimate	0.68	0.62	0.62	0.58	0.61	0.57	0.52	0.60	0.57	0.52	0.21	0.32	0.54
Backcast	0.38	0.34	0.36	0.33	0.33	0.31	0.29	0.34	0.34	0.30	-	-	0.28
MF-FAVAR: no VAT data													
Nowcast	1.09	0.95*	1.04	0.97	0.99	0.90	0.84	0.94	0.95	0.87	0.72	0.70	0.91
Estimate	0.61	0.55	0.58	0.55	0.57	0.50	0.46	0.53	0.52	0.47	0.21	0.31	0.49
Backcast	0.33	0.30	0.34	0.32	0.31	0.27	0.25	0.30	0.31	0.27	-	-	0.25
MF-FAVAR: with VAT data													
Nowcast	1.05*	0.98	1.03	1.06	1.15	1.02	0.85*	1.02	1.01	0.83*	0.64	0.64*	0.94
Estimate	0.60*	0.58	0.61	0.62	0.64	0.58	0.49*	0.59	0.59	0.45*	0.21	0.29	0.52
Backcast	0.33	0.33	0.35	0.35	0.34	0.32	0.27	0.34	0.34	0.25	-	-	0.27
MF-FAVAR: with VAT data and lag=1 month													
Nowcast	1.05*	0.98	1.04	1.06	1.15	1.02	0.85*	1.02	1.01	0.83*	0.64	0.64*	0.94
Estimate	0.60*	0.58	0.61	0.62	0.64	0.58	0.49*	0.59	0.59	0.45*	0.21	0.29	0.52
Backcast	0.33	0.33	0.35	0.35	0.34	0.32	0.27	0.34	0.34	0.25	-	-	0.27

Notes: * denotes rejection of the null of equal forecast accuracy against the benchmark MF-VAR model at the 0.10 significance level using a two-sided Diebold and Mariano (1995) test. All models do not have the cross-sectional restrictions imposed. All factors are estimated using the TP approach.